

CTP # 2213

On Coarse-grained Entropy and Stimulated Emission in Curved Space-Time

Esko Keski-Vakkuri*

Center for Theoretical Physics

Laboratory for Nuclear Science

and Department of Physics

Massachusetts Institute of Technology

77 Massachusetts Avenue, Cambridge MA 02139

August, 1993

Abstract

We study the entropy generation and particle production in scalar quantum field theory in expanding spacetimes with many-particle mixed initial states. The recently proposed coarse-grained entropy approach by Brandenberger *et al.* is applied to systems which may have a non-zero initial entropy. We find that although the particle production is amplified as a result of boson statistics, the (coarse-grained) entropy generation is *attenuated* when initial particles are present.

*Supported in part by funds provided by the U.S. Department of Energy (DOE) under contract #DE-AC02-76ER03069.

One of the interesting features of quantized fields in a curved spacetime [1] is that the concept of particles becomes very observer-dependent. For instance, in an expanding Universe spontaneous particle creation can occur. One defines generally a vacuum state such that all inertial observers in the past region agree that the spacetime looks empty of particles. As a result of the expansion of the Universe, the above vacuum state looks full of particles using modes natural to inertial observers in the far future region. Stated differently, a no-particle initial state can evolve to a many-particle state. However, since one starts with a pure state, one ends with a pure state. Thus there must be subtle correlations between the particles in the final state. In particular, there is no entropy production in this process even if lots of particles are produced. But, it may be that some of these subtle correlations are very difficult to detect and/or that they may be quite sensitive to interactions between the produced particles. One may then contemplate a picture where such information about the system is considered to be “less relevant” and either discarded altogether or some kind of “statistical averaging” procedure is applied to it. This way one can try to associate a “coarse-grained” entropy to the final state of the system, hopefully in as natural way as possible. There has been a lot of work in this direction by Hu, Kandrup and collaborators [2].

Recently, two different such approaches have been proposed. Brandenberger, Mukhanov and Prokopec (BMP) discussed in [3, 4] a coarse-graining procedure based on averaging over the so called squeeze angles which appear in the S-matrix of particle production. On the other hand, Gasperini and Giovannini [5], together with Veneziano (GGV) [6] related entropy generation to an increased dispersion of a superfluctuant operator. Both groups were especially interested in the entropy generation related to the production of gravitational waves and density fluctuations in inflationary universe models.

In this work, our aim is to follow the BMP approach and study the entropy generation starting not from an initial vacuum state with zero entropy, but allowing the system to be initially in some generic many-particle (mixed) state with non-zero entropy. If one starts with many bosons it is known [7] that the particle production will be amplified as a result of boson statistics, as one would expect. So, in general one can ask whether the entropy generation (in the coarse-grained sense) would also be amplified or not. Indeed, as a consistency check it is necessary to investigate if definitions of coarse-grained entropy will lead to a growing entropy even if initial state is allowed to be an arbitrary many-particle state with initial entropy. In [5, 6] the GGV entropy was shown to be growing at least in certain classes of initial states. Interestingly, it was found that their entropy generation did not depend at all on the number of particles or entropy of the initial state. Here we will attempt to investigate the BMP entropy in similar situations. At least in the case of an initial density matrix where particles appear as pairs of opposite momenta, and initial entropy depends on the average occupation number per mode, we can show that the BMP entropy grows, though the entropy generation is *attenuated*. The BMP entropy *does* depend on the initial number of particles in a non-trivial way. In the end we comment briefly on the case of an initial thermal density matrix.

A scalar field in a D -dimensional curved spacetime is described by an action

$$S = \int d^D x \sqrt{-g} \frac{1}{2} [g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - (m^2 + \xi R(x)) \phi^2], \quad (1)$$

where $R(x)$ is the Ricci scalar curvature of the metric and ξ is a coupling constant. Assume

that the metric depends explicitly on time and that it is asymptotically flat in the far past and far future [1]:

$$g_{\mu\nu}(\vec{x}, t) \rightarrow C_{\pm} \eta_{\mu\nu} \quad (2)$$

as $t \rightarrow \pm\infty$. In this case there are two natural ways to quantize the field ϕ in Heisenberg picture :

$$\phi = \sum_{\vec{k}} a_{\vec{k}}^{in} u_{\vec{k}}^{in} + h.c. \quad (3)$$

$$\phi = \sum_{\vec{k}} a_{\vec{k}}^{out} u_{\vec{k}}^{out} + h.c. \quad (4)$$

where $u_{\vec{k}}^{in}$ looks like a plane wave in the far past,

$$u_{\vec{k}}^{in}(\vec{x}, t) \propto e^{-i\omega_{\vec{k}}^{in}t + i\vec{k}\cdot\vec{x}}, \quad t \rightarrow -\infty \quad (5)$$

and $u_{\vec{k}}^{out}$ looks like a plane wave in the far future, respectively

$$u_{\vec{k}}^{out}(\vec{x}, t) \propto e^{-i\omega_{\vec{k}}^{out}t + i\vec{k}\cdot\vec{x}}, \quad t \rightarrow +\infty. \quad (6)$$

These solutions can be related via a Bogoliubov transformation, which can be given in terms of annihilation/creation operators as

$$a_{\vec{k}}^{in} = \alpha_{\vec{k}\vec{p}}^* a_{\vec{p}}^{out} - \beta_{\vec{k}\vec{p}}^* a_{\vec{p}}^{\dagger out}. \quad (7)$$

This transformation is generated by a S-matrix

$$a_{\vec{k}}^{in} = \mathbf{S} a_{\vec{k}}^{out} \mathbf{S}^{-1}, \quad (8)$$

which has the explicit form [8]

$$\mathbf{S} = \frac{1}{\sqrt{\det(\alpha_{\vec{k}\vec{p}})}} : \exp\left\{\frac{1}{2}[\alpha^{-1}\beta^*]_{\vec{k}\vec{k}'} a_{\vec{k}}^{\dagger out} a_{\vec{k}'}^{out} + [\alpha^{-1} - 1]_{\vec{k}\vec{k}'} a_{\vec{k}}^{\dagger out} a_{\vec{k}'}^{out} - \frac{1}{2}[\alpha^{-1}\beta]_{\vec{k}\vec{k}'} a_{\vec{k}}^{out} a_{\vec{k}'}^{out}\right\} : \quad (9)$$

The factor $\frac{1}{\sqrt{\det(\alpha_{\vec{k}\vec{p}})}}$ is the in-out vacuum amplitude. We use the convention of [9] where the coefficients α have taken to be real. The S-matrix is known to generate a unitary transformation between the 'in' and 'out' representations if the gravitational field has a compact support [10]. For Robertson-Walker type universes the in-out vacuum amplitude is zero and the 'in' and 'out' representations are thus unitarily inequivalent.

The S-matrix relates the in- and out-vacuum states in the following way

$$\begin{aligned} |0, in\rangle &= \mathbf{S} |0, out\rangle \\ &= \frac{1}{\sqrt{\det(\alpha)}} \exp\left\{\frac{1}{2}[\alpha^{-1}\beta]_{\vec{k}\vec{k}'} a_{\vec{k}}^{\dagger out} a_{\vec{k}'}^{out}\right\} |0, out\rangle. \end{aligned} \quad (10)$$

This is the statement that an inertial observer in the far future region sees the in-vacuum state as full of out-particles. Similarly, the density matrix of the system expanded using in-modes ($\equiv \rho_i$) can be related to an expression using out-modes ($\equiv \rho_f$) as follows

$$\begin{aligned}
\rho_i &= \prod_{\vec{k}} \sum_{n_{\vec{k}}=0}^{\infty} f(n_{\vec{k}}) |n_{\vec{k}}, in\rangle \langle in, n_{\vec{k}}| \\
&= \prod_{\vec{k}} \sum_{n_{\vec{k}}} \frac{f(n_{\vec{k}})}{n_{\vec{k}}!} \mathbf{S} a_{\vec{k}}^{\dagger out} \mathbf{S}^{-1} \dots \mathbf{S} a_{\vec{k}}^{\dagger out} \mathbf{S}^{-1} \mathbf{S} |0, out\rangle \langle out, 0| \mathbf{S}^{-1} \mathbf{S} a_{\vec{k}}^{out} \mathbf{S}^{-1} \dots \mathbf{S} a_{\vec{k}}^{out} \mathbf{S}^{-1} \\
&= \prod_{\vec{k}} \sum_{n_{\vec{k}}=0}^{\infty} f(n_{\vec{k}}) \mathbf{S} |n_{\vec{k}}, out\rangle \langle out, n_{\vec{k}}| \mathbf{S}^{-1} \equiv \mathbf{S} \rho_f \mathbf{S}^{-1}.
\end{aligned} \tag{11}$$

Suppose now that the system is initially in an arbitrary many-particle state. In this state the average occupation number per mode (using in-modes) is given by

$$\bar{n}_{\vec{k}}^i = \frac{1}{Tr \rho_i} Tr(\rho_i a_{\vec{k}}^{\dagger in} a_{\vec{k}}^{in}). \tag{12}$$

In the far future region an inertial observer sees the average occupation number per mode using out-modes as

$$\bar{n}_{\vec{k}}^f = \frac{1}{Tr \rho_f} Tr(\mathbf{S} \rho_f \mathbf{S}^{-1} a_{\vec{k}}^{\dagger out} a_{\vec{k}}^{out}) \tag{13}$$

Using the cyclicity of the trace and the properties of the Bogoliubov transformation one can derive the relation between $\bar{n}_{\vec{k}}^f$ and $\bar{n}_{\vec{k}}^i$ to be

$$\bar{n}_{\vec{k}}^f = \bar{n}_{\vec{k}}^i + |\beta_{\vec{k}\vec{p}}|^2 (1 + 2\bar{n}_{\vec{p}}^i). \tag{14}$$

This is the formula for “stimulated emission” [7]. It tells us that even if the spontaneous creation of particles is weak, $|\beta_{\vec{k}\vec{p}}|^2 \ll 1$, the particle production can become arbitrarily large,

$$\Delta \bar{n}_{\vec{k}} \equiv \bar{n}_{\vec{k}}^f - \bar{n}_{\vec{k}}^i = |\beta_{\vec{k}\vec{p}}|^2 (1 + 2\bar{n}_{\vec{p}}^i) \rightarrow large \tag{15}$$

if the initial average occupation number per mode is arbitrarily large. This amplification of particle production is a result of the boson statistics of the particles. For fermions the particle production would be attenuated [11].

Let us now discuss for notational simplicity metrics of the form

$$ds^2 = dt^2 - a^2(t) d^2 \vec{x}^2, \tag{16}$$

where $a(t)$ is a scale parameter of the universe. We again just require that $a(t) \rightarrow a_{\pm}$ asymptotically as $t \rightarrow \pm\infty$. As a result of the invariance under spatial translations, the Bogoliubov coefficients can be written as

$$\begin{aligned}
\alpha_{\vec{k}\vec{p}} &= \alpha_{\vec{k}} \delta_{\vec{k},\vec{p}} \equiv \cosh r_{\vec{k}} \delta_{\vec{k},\vec{p}} \\
\beta_{\vec{k}\vec{p}} &= \beta_{\vec{k}} \delta_{\vec{k},-\vec{p}} \equiv e^{i\theta_{\vec{k}}} \sinh r_{\vec{k}} \delta_{\vec{k},-\vec{p}}.
\end{aligned} \tag{17}$$

The parameters $r_{\vec{k}}$, $\theta_{\vec{k}}$ are called squeeze parameter and squeeze angle in the Quantum Optics literature, and the S-matrix is called a two-mode squeeze operator [12]. If one starts with

a vacuum state, the final state (10) is called a squeezed vacuum. In the initial vacuum case, if one expands the corresponding $\mathbf{S}\rho_f\mathbf{S}^{-1}$ in the 'out' basis of energy eigenstates, one finds [4] that the off-diagonal components of $\mathbf{S}\rho_f\mathbf{S}^{-1}$ have an oscillatory dependence of the angles θ_k . In the BMP coarse-grained entropy approach it is assumed that these angles represent irrelevant information about the system (*e.g.*, in the sense that they would be very difficult to measure) and they are therefore averaged over. After the averaging only the diagonal elements of $\mathbf{S}\rho_f\mathbf{S}^{-1}$ survive and one then defines a coarse-grained entropy with the resulting reduced density matrix ρ_{red} with the usual formula $S = -k_B \text{Tr}(\rho_{red} \log \rho_{red})$. The result is [3, 4]

$$\begin{aligned} S^{f,0} &\equiv \sum_{\vec{k}} s_{\vec{k}}^{f,0} \equiv k_B \sum_{\vec{k}} (\cosh^2 r_{\vec{k}} \log \cosh^2 r_{\vec{k}} - \sinh^2 r_{\vec{k}} \log \sinh^2 r_{\vec{k}}) \\ &= k_B \sum_{\vec{k}} [(\bar{n}_{\vec{k}}^{f,0} + 1) \log(\bar{n}_{\vec{k}}^{f,0} + 1) - (\bar{n}_{\vec{k}}^{f,0}) \log(\bar{n}_{\vec{k}}^{f,0})] , \end{aligned} \quad (18)$$

where the notation $\bar{n}_{\vec{k}}^{f,0}$ means the LHS of (14) in the case of an initial vacuum state. Now we turn to consider initial density matrices ρ_i which can describe generic many-particle states with non-zero entropy. Let us assume that the initial density matrix has the form

$$\rho_i = \prod_{\vec{k}, (k_z > 0)} \sum_{n=0}^{\infty} f_{\vec{k}}(n) |n_{\vec{k}}, n_{-\vec{k}}, in\rangle \langle in, n_{\vec{k}}, n_{-\vec{k}}| , \quad (19)$$

where the coefficients $f(n_{\vec{k}})$ are of the form $f(n_{\vec{k}}) = (\bar{n}_{\vec{k}}^i)^n / (\bar{n}_{\vec{k}}^i + 1)^{n+1}$. That is, we start with a many-particle state where particles appear in pairs of opposite momenta, with an initial average occupation number spectrum $\bar{n}_{\vec{k}}^i = \bar{n}_{-\vec{k}}^i$ and with (ordinary) entropy given by $-k_B \text{Tr}(\rho_i \log \rho_i)$. Writing the initial entropy in more explicit form, it is

$$S^i \equiv \sum_{\vec{k}, (k_z > 0)} s_{\vec{k}}^i = k_B \sum_{\vec{k}, (k_z > 0)} [(\bar{n}_{\vec{k}}^i + 1) \log(\bar{n}_{\vec{k}}^i + 1) - (\bar{n}_{\vec{k}}^i) \log(\bar{n}_{\vec{k}}^i)] . \quad (20)$$

When we expand the resulting final density matrix $\mathbf{S}\rho_f\mathbf{S}^{-1}$ (11) in 'out' energy eigenstates, we find that also in this case the off-diagonal elements have an oscillatory dependence on the angles $\theta_{\vec{k}}$. Therefore, following the BMP approach and averaging over the angles, only the diagonal elements will survive. Thus the reduced density matrix of the final state has the form

$$\rho_{red} = \prod_{\vec{k}, (k_z > 0)} \sum_{n=0}^{\infty} \bar{f}_{\vec{k}}(n) |n_{\vec{k}}, n_{-\vec{k}}, out\rangle \langle out, n_{\vec{k}}, n_{-\vec{k}}| , \quad (21)$$

where $\bar{f}_{\vec{k}}(n) = \langle n_{\vec{k}}, n_{-\vec{k}}, out | \mathbf{S}\rho_f\mathbf{S}^{-1} | out, n_{\vec{k}}, n_{-\vec{k}} \rangle$. After some effort, one can show that the coefficients have the form

$$\bar{f}_{\vec{k}}(n) = \frac{(\bar{n}_{\vec{k}}^f)^n}{(\bar{n}_{\vec{k}}^f + 1)^{n+1}} , \quad (22)$$

where $\bar{n}_{\vec{k}}^f$ is the LHS of (14) with the $\bar{n}_{\vec{k}}^i$ of (19). Thus, the final coarse-grained entropy is

$$S^f \equiv \sum_{\vec{k}, (k_z > 0)} s_{\vec{k}}^f = k_B \sum_{\vec{k}, (k_z > 0)} [(\bar{n}_{\vec{k}}^f + 1) \log(\bar{n}_{\vec{k}}^f + 1) - (\bar{n}_{\vec{k}}^f) \log(\bar{n}_{\vec{k}}^f)] . \quad (23)$$

The entropy depends only on the occupation number spectrum of particles in the final state. This result is in agreement with a similar formula given in [3] by a more heuristic argument to define entropy of a statistical system with a definite spectrum which is valid both in and far out of thermodynamical equilibrium. Let us now compare the entropy generation per mode in the initial vacuum and initial many-particle cases. Denote

$$\Delta_0 s_{\vec{k}} \equiv s_{\vec{k}}^{f,0} - 0 \quad (24)$$

$$\Delta s_{\vec{k}} \equiv s_{\vec{k}}^f - s_{\vec{k}}^i, \quad (25)$$

where (24) applies to the former case and (25) to the latter case. As a first consistency check, we find that

$$\Delta s_{\vec{k}} \geq 0 \quad (26)$$

so that the coarse-graining led to a growing entropy in our many-particle case. However, as we compare (25) and (24) we find that

$$\Delta s_{\vec{k}} \leq \Delta_0 s_{\vec{k}}; \quad (27)$$

i.e., the entropy generation is *attenuated* if one starts with many particles present in the mode $\vec{k}$. The equality holds iff $\bar{n}_{\vec{k}}^i = 0$. The difference $\Delta s_{\vec{k}} - \Delta_0 s_{\vec{k}}$ depends symmetrically on $\bar{n}_{\vec{k}}^i$ and $|\beta_{\vec{k}}|^2$ and decreases monotonically as either variable increases. Further, as both variables approach infinity,

$$\Delta s_{\vec{k}} - \Delta_0 s_{\vec{k}} \rightarrow \log 2 - 1 \approx -0.31 \quad (28)$$

asymptotically. This finite value is the maximum difference between the generated entropies per mode. Thus, unlike the GGV entropy, the BMP entropy is *not* independent of the number of particles in the initial state, but has some 'memory' about the initial occupation numbers. Since entropy is a measure of loss of information, it would appear that more information about the initial state of the system is conveyed to the final coarse-grained state when stimulated emission dominates the spontaneous particle production (since entropy generation is attenuated).

Next case to be investigated would be an initial thermal density matrix

$$\rho_i = \prod_{\vec{k}} Z_{\vec{k}}^{-1} \exp(-\beta \omega_{\vec{k}}^{in} a_{\vec{k}}^{\dagger in} a_{\vec{k}}^{in}).$$

This situation is somewhat trickier to deal with, for the following reason. Initially, the particles of opposite momenta are uncorrelated. However, in the expansion of the universe the particles are produced in pairs of opposite momenta. This induces correlations between the opposite momenta in the final density matrix. It would have the form

$$\mathbf{S} \rho_f \mathbf{S}^{-1} = \prod_{\vec{k}, (k_z > 0)} \sum_{n, m, n', m'=0}^{\infty} f_{\vec{k}}(n, m; n', m') | n_{\vec{k}}, m_{-\vec{k}}, out \rangle \langle out, n'_{\vec{k}}, m'_{-\vec{k}} |, \quad (29)$$

where

$$f_{\vec{k}}(n, m; n', m') = \langle n_{\vec{k}}, m_{-\vec{k}}, out | \frac{1}{Z_{\vec{k}} Z_{-\vec{k}}} e^{-\beta \omega_{\vec{k}}^{in} [(\alpha_{\vec{k}} a_{\vec{k}}^{\dagger out} - \beta_{\vec{k}} a_{-\vec{k}}^{out})(\alpha_{\vec{k}}^* a_{\vec{k}}^{out} - \beta_{\vec{k}}^* a_{-\vec{k}}^{\dagger out}) + (\vec{k} \leftrightarrow -\vec{k})]} | out, n'_{\vec{k}}, m'_{-\vec{k}} \rangle. \quad (30)$$

Again, one can see that the $n \neq n'$ or $m \neq m'$ components have an oscillatory dependence of the squeeze angles and they vanish in the coarse graining. However, the diagonal coefficients (those of the reduced density matrix) will have a form $f_{\vec{k}}(n, m)$ where the dependence on n and m does not factorize. Hence the opposite momenta have acquired correlations through the particle production and the reduced density matrix is not of the same type as the initial one. As advocated in [6], one would like to ignore the correlations between different modes. In [6] this problem was avoided by replacing the two-mode squeeze operator by a one-mode squeeze operator, but this does not appear entirely satisfactory, since the particles are then not created in the correct way as pairs of opposite momenta. We would like to propose that the correlations between opposite momenta could be ignored by proceeding to define $\bar{f}_{\vec{k}}(n) = \sum_m f_{\vec{k}}(n, m)$ and $\bar{f}_{-\vec{k}}(m) = \sum_n f_{\vec{k}}(n, m)$. Then we would define the final reduced density matrix to be

$$\rho_{red} = \prod_{\vec{k}, (k_z > 0)} \sum_{n, m=0}^{\infty} \bar{f}_{\vec{k}}(n) \bar{f}_{-\vec{k}}(m) |n_{\vec{k}}, m_{-\vec{k}}, out\rangle \langle out, n_{\vec{k}}, m_{-\vec{k}}|, \quad (31)$$

which is of the same type as the initial density matrix. Now the final entropy would be given by

$$S^f = -k_B \sum_{\vec{k}} \sum_n \bar{f}_{\vec{k}}(n) \log \bar{f}_{\vec{k}}(n). \quad (32)$$

Unfortunately, at the present we do not have explicit formulas for the coefficients $\bar{f}_{\vec{k}}(n)$ or the final entropy. It would be very interesting to see if the resulting expressions could depend on the final average occupation number in the same fashion as in the earlier case. We hope to be able to return to this question in the future.

Finally, let us clarify that even if we found a different result as in the GGV approach, namely an entropy that depends on the number of particles in the initial state, we are not arguing that it would mean that the BMP approach is 'better' than the GGV approach. As stated in [2], it is good to have different definitions of entropy, corresponding to loss of different information about the system. Both BMP and GGV approaches have the virtue of giving the correct average occupation number for particles in the final state. Otherwise the GGV approach appears to discard information about the system a bit more generously, since it leads to a greater growth of entropy.

Acknowledgements

It is a pleasure to thank prof. S.D. Mathur for guidance and numerous discussions. I am also grateful to profs. R. Brandenberger, A.J. Niemi and S. Stenholm for helpful discussions.

References

- [1] N.D. Birrell and P.C.W. Davies, *Quantum Fields in Curved Space*, Cambridge Univ. Press (1982)
- S. Fulling, *Aspects of Quantum Field Theory in Curved Space-Time*, Cambridge Univ. Press (1989)

- [2] B. Hu and D. Pavon, Phys. Lett. **B180**, 329 (1986)
H. Kandrup, J. Math. Phys. **28**, 1398 (1987); Phys. Rev. **D37**, 3505 (1988)
B. Hu and H. Kandrup, Phys. Rev. **D35**, 1776 (1987)
S. Habib and H. Kandrup, Ann. Phys. **191**, 335 (1989)
- [3] R. Brandenberger, V. Mukhanov and T. Prokopec, *The Entropy of the Gravitational Field*, Brown Univ. preprint BROWN-HET-849 (1992); Phys. Rev. Lett. **69**, 3606 (1992)
- [4] T. Prokopec, *Entropy of the Squeezed Vacuum*, Brown Univ. preprint BROWN-HET-861 (1992)
- [5] M. Gasperini and M. Giovannini, Phys. Lett. **B301**, 334 (1993); *Quantum Squeezing and Cosmological Entropy Production*, CERN preprint CERN-TH.6954/93, gr-qc@xxx/9307024
- [6] M. Gasperini, M. Giovannini and G. Veneziano, Phys. Rev. **D48**, R439 (1993)
- [7] L. Parker, Phys. Rev. **183**, 1057 (1969)
R.M. Wald, Phys. Rev. **D13**, 3176 (1976)
J. Audretsch, in *Quantum Mechanics in Curved Space-Time*, ed. J. Audretsch and V. de Sabbata, Plenum Press (1990)
- [8] The explicit formula for the S-matrix can be found (in the case of a RW-universe) *e.g.* in J. Ambjørn, R.J. Hughes and N.K. Nielsen, Ann. Phys. **150**, 92 (1983). Note that there are minor differences in conventions used in that paper and in this work.
- [9] B. DeWitt, Phys. Rep. **19**, 297 (1975)
- [10] R.M. Wald, Ann. Phys. **118**, 490 (1979)
- [11] L. Parker, Phys. Rev. **D3**, 346 (1971)
- [12] B. Schumaker, Phys. Rep. **135**, 317 (1986)